

FRACTURE PROPAGATION: Recent Research and Implications

Story by Karl Birkeland, Jürg Schweizer, and Bruce Jamieson


A quick perusal through some recent ISSW Proceedings demonstrates a greater emphasis on fracture propagation. The Whistler ISSW boasted 16 papers and posters on fracture propagation and associated field tests, and the evening's beer-fueled discussions often came around to talk of fracture propagation. In her usual nice way, TAR editor Lynne Wolfe cornered us at that conference and encouraged us to write a quick article summarizing some of this work; out of her encouragement came this article. We attempt to synthesize some of the recent research and ideas, and discuss some of the implications for avalanche practitioners.

Propagation versus Initiation

While several new tests are available, many of the ideas about fracture propagation are not new. We have known implicitly that avalanches require both fracture initiation and propagation, though this has only

been emphasized in our writing and teaching for about the last five or 10 years (e.g., Schweizer, et al., 2003). Until quite recently, our field tests and much of the research have predominantly emphasized initiation over propagation. Likewise, some of the recent new models of fracture propagation simply provide an improved explanation for phenomena that practitioners have known and observed for many decades, such as avalanches remotely triggered from flat terrain.

This most recent emphasis on fracture propagation began when researchers formalized what many practitioners had long observed about stability tests – that it is important to not only observe how much dynamic force (or load) it takes to fracture a weak layer, but also to look at how that fracture occurs. With rutschblocks this meant observing the type of release (whole block, part of the block, or the edge), as noted by Schweizer, et al., (1995a).



An excavated whumpf shows the fractured and collapsed weak layer on the left and the intact weak layer on the right.
Photo by University of Calgary ASARC


A later survey of avalanche forecasters indicated they were typically more interested in the way a rutschblock fractured than in the actual rutschblock score (Schweizer and Wiesinger, 2001). With compression tests and stuffblock tests, researchers began to look at shear quality (Birkeland and Johnson, 1999; Johnson and Birkeland, 2002) and fracture character (van Herwijnen and Jamieson, 2002, 2004). Johnson and Birkeland (2002) hypothesized that shear quality

might provide a qualitative measure (at a small scale) of how a fracture might propagate. Observing the way these tests fractured and whether fractures were “clean” or “sudden” had been done by practitioners for a long time, but research showed that taking this into account when interpreting stability tests could reduce the number of “false stables” or test results indicating stable conditions on slopes that showed other obvious signs of instability (Johnson and Birkeland, 2002; Birkeland and Chabot, 2006; Schweizer et al., 2006).

New tests focusing specifically on fracture propagation expanded on observations of shear quality, fracture character, and rutschblock release type. Gauthier and Jamieson (2006; 2008a), Sigrist (2006), and Sigrist and Schweizer (2007) came up with a test that involved isolating a column parallel to the fall line of the slope, initiating a fracture with a saw, and observing whether or not the fracture then self-propagated through the rest of the column. This test was refined and eventually dubbed the Propagation Saw Test (PST). Simenhois and Birkeland (2006) simultaneously and independently worked on the Extended Column Test (ECT), in which a 90cm-wide cross-slope column is loaded on one side with loading steps identical to the compression test.

Both the PST and ECT quickly found their way into the toolboxes of many practitioners, with recent research showing the ECT to have a low false-stability rate and the PST to have a low false-instability rate. Since there are already several articles on both of these tests, we won't go into further details here (in addition to the papers already cited, check out Birkeland and Simenhois, 2008; Gauthier and Jamieson, 2008b; Moner et al., 2008; Ross and Jamieson, 2008; Winkler and Schweizer, 2008). One advantage of the new tests is that we can now begin to investigate some of the factors affecting fracture propagation in the field, such as changes in slab depth (Simenhois and Birkeland, 2008a) or surface warming (Simenhois and Birkeland, 2008b).

In addition to some of the new tests focused on fracture propagation, there has been renewed interest in the theory behind fracture propagation. Johnson et al., (2004) measured the speed of a propagating fracture across a flat meadow utilizing geophones, finding that the fracture traveled at 20 ± 2 m s⁻¹. van Herwijnen and Jamieson (2005) measured fracture speeds with a high speed camera, calculating speeds between 17 and 26 m s⁻¹. These fracture speed measurements helped to reignite a mostly dormant



For practitioners, we need to realize that determining instability requires understanding and assessing both initiation and propagation.

Jürg Schweizer triggers a whumpf resulting in a surface crack on a slope near Davos. Photo by SLF

debate about the relative importance of collapse in fracture propagation. The speeds are consistent with theory proposing collapse as a driving force of fracture propagation (Johnson, 2001; van Herwijnen and Jamieson, 2005), though other work suggests they are also consistent with existing models of shear fracture propagation (McClung, 2005). Gauthier and Jamieson (2008a) reported similar PST results on slopes as on adjacent flat terrain, an observation that supports the importance of collapse in driving fracture propagation.

High-speed videos show a variety of weak layers – including thin weak layers – collapsing and shearing, with none of them showing slope-parallel shear fracture without collapse (Schweizer et al., 1995b; van Herwijnen and Jamieson, 2005; van Herwijnen et al., 2008).

Of course, the idea that collapse plays a role in fracture propagation is not new. After all, snow in general and weak layers in particular are highly porous, making collapse possible. Seligman (1936) noted that avalanches could be triggered from flat terrain, and Bader (1951) stated collapse as one of several methods for fracture under the slab. Bradley (1968) developed a field instrument and method for forecasting avalanches related to collapse in depth hoar snowpacks by calculating a bulk strength-to-load index. Lackinger (1989) described the bending wave due to a collapsing weak layer. Johnson (2001) wrote a simple equation for the bending wave and better described remote triggering from low-angle terrain. Heierli and colleagues (Heierli, 2005; Heierli et al., 2008; Heierli and Zaiser, 2006; Heierli and Zaiser, 2008) greatly improved the mathematical description for the collapse and bending wave and then combined this theory with shear-fracture theory.

In essence, this most recent work allows for both collapse and shear as potential driving forces behind fracture propagation (Sigrist, 2006; Heierli et al., 2008). Fractures in thinner weak layers and on steeper slopes are predicted to be more dependent on shear, while fractures in thicker weak layers and on flatter terrain are more dependent on collapse.

What Does this Mean to Us?

While plenty of new information is available for pondering the theory behind fracture propagation, the big question is: “What are the implications for practitioners?” Before we address this question, we need to remember that the basic observations we have made for years are still valid. For example, practitioners have known of and observed avalanche triggering from flat terrain for many decades. Some of the new theory simply gives us a better mathematical description for that observation. Further, we have known – at least intuitively – that both fracture initiation and fracture propagation are necessary for avalanches.

The first implication for practitioners of some of the new work is that researchers and practitioners need to consider both the slab and the weak layer. Our emphasis in the past has been on fracture initiation, and we tended to focus primarily on the weak layer. This shifted as we came to better understand the role of the slab in initiation, and as we now start to gain knowledge of the role of the slab in



Tests in flat areas might be useful for predicting conditions on nearby slopes.

On this part of a spring tour in the Rockies, Dave Gauthier (in background), Antonia Zeidler (foreground), and Bruce Jamieson were triggering whumpfs every few steps.
Photo by Bruce Jamieson

propagation we are realizing that it is vitally important to look at both the slab and weak layer together to better understand avalanches. Practitioners should be sure to note the characteristics of when fractures are propagating and integrate this knowledge into stability assessments.

A second implication for practitioners is that we need to realize that determining instability requires understanding and assessing both initiation and propagation. Luckily, we now have – besides things like the rutschblock release type, fracture character, and shear quality – two tests (the PST and ECT) that give us a start at specifically indexing fracture propagation potential, thereby providing us with new methods for assessing snow stability. We also need to be better aware of how fracture-propagation propensity might vary spatially around starting zones. This is an open question, though some preliminary work has been done (Birkeland and Simenhois, 2008; Hendrikx and Birkeland, 2008; Hendrikx et al., in press).

A third implication has to do with the location of our field tests. We don't know if fractures are occurring first in shear or in compression. However, it is clear that both shear and collapse are occurring in some sort of mixed mode and that collapse is an essential energy source for propagation in some cases. Thus, for collapsible weak layers, tests in flat areas or in areas with shallow slope angles might be useful for predicting conditions on nearby slopes, as long as the snow stratigraphy in those flat areas is representative of the slopes in question.

Indeed, a limited dataset shows this to be the case for the PST (Gauthier and Jamieson, 2008a), and some limited and preliminary data from this season suggest that ECTs in flat terrain may provide useful information about the potential for remote triggering (Simenhois, pers. comm., 2009). If these findings are confirmed, it would be extremely helpful for practitioners and recreationists since safe pit sites would be much easier to locate in the flats or at least on gentle slopes than on steep slopes during unstable conditions. Of course, some layers (such as poorly bonded crusts or weak interfaces) are much less collapsible. If they don't

collapse – and so far, high-speed videos have not revealed any that don't have at least some collapse – then tests involving such weak interfaces will need to be conducted on slopes.

New Ways to Look at Snowpack

Recent research on fracture propagation and the development of tests attempting to index fracture-propagation propensity provide all of us with new tools and new ways to look at the snowpack.

It is important to remember that these new tools and insights don't replace our proven tools developed over the past several decades, but instead they simply add to our toolbox. Avalanche forecasting and stability assessment still require a holistic approach that takes into account diverse data including weather, avalanche, and snowpack observations. The key is to look into the snow, poke around, and do a variety of tests, while at the same time realizing that there is no ultimate test. Hopefully future research, combined with careful observations by practitioners, will continue to improve our understanding and our methods for evaluating slope stability.

REFERENCES

- Bader, H., B.L. Hansen, J.A. Joseph and M.A. Sandgren, 1951. Preliminary investigations of some physical properties of snow. *SIPRE Report 7*: 48 pp.
- Birkeland, K.W. and Chabot, D., 2006. *Minimizing “false-stable” stability test results: Why digging more snowpits is a good idea*. In: A. Gleason (Editor), Proceedings of the 2006 International Snow Science Workshop, Telluride, Colorado, pp. 498-504.
- Birkeland, K.W. and Johnson, R.F., 1999. The stuffblock snow stability test: comparability with the rutschblock, usefulness in different snow climates, and repeatability between observers. *Cold Regions Science and Technology*, 30(1-3): 115-123.
- Birkeland, K.W. and Simenhois, R., 2008. *The extended column test: Test effectiveness, spatial variability, and comparison with the propagation saw test*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 401-407.
- Bradley, C.C., 1968. The resistograph and the compressive strength of snow. *Journal of Glaciology*, 7(51): 499-506.
- Gauthier, D. and Jamieson, J.B., 2006. Towards a field test for fracture propagation propensity in weak snowpack layers. *Journal of Glaciology*, 52(176): 164-168.
- Gauthier, D. and Jamieson, J.B., 2008a. Fracture propagation propensity in relation to snow slab avalanche release: Validating the propagation saw test. *Geophysical Research Letters*, 35(L13501): doi: 10.1029/2008GL034245.
- Gauthier, D. and Jamieson, J.B., 2008b. *Predictions of the propagation saw test: Comparisons with the other instability tests at skier tested slopes*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 408-414.
- Heierli, J., 2005. Solitary fracture waves in metastable snow stratifications. *Journal of Geophysical Research*, 110: doi: 10.1029/2004JF000178.
- Heierli, J., Gumbsch, P. and Zaiser, M., 2008. Anticrack nucleation as triggering mechanism for snow slab avalanches. *Science*, 321: 240-243.
- Heierli, J. and Zaiser, M., 2006. An analytical model for fracture nucleation in collapsible stratifications. *Geophysical Research Letters*, 33: L06501, doi: 10.1029/2005GL025311.
- Heierli, J. and Zaiser, M., 2008. Failure initiation in snow stratifications containing weak layers: Nucleation of whumpfs and slab avalanches. *Cold Regions Science and Technology*: doi: 10.1016/j.coldregions.2007.02.07.
- Hendrikx, J. and Birkeland, K.W., 2008. *Slope scale spatial variability across time and space: Comparison of results from continental and maritime climates*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, pp. 155-162.
- Hendrikx, J., Birkeland, K.W. and Clark, M., In press. Assessing changes in the spatial variability of the snowpack fracture propagation propensity over time. *Cold Regions Science and Technology*: doi: 10.1016/j.coldregions.2008.12.001.
- Johnson, B.C., 2001. *Remotely triggered slab avalanches*. MSc Thesis, University of Calgary, Calgary AB, Canada, 98 pp.
- Johnson, B.C., Jamieson, J.B. and Stewart, R.R., 2004. Seismic measurement of fracture speed in a weak snowpack layer. *Cold Regions Science and Technology*, 40(1-2): 41-46.
- Johnson, R.F. and Birkeland, K.W., 2002. *Integrating shear quality into stability test results*. In: J.R. Stevens (Editor), International Snow Science Workshop, Penticton, B.C., pp. 508-513.
- Lackinger, B., 1989. Supporting forces and stability of snow-slab avalanches: a parameter study. *Annals of Glaciology*, 13: 140-145.
- McClung, D.M., 2005. Approximate estimates of fracture speeds for dry slab avalanches. *Geophysical Research Letters*, 32: doi: 10.1029/2005GL022391.
- Moner, I., Gavalda, J., Bacardit, M., Garcia, C. and Marti, G., 2008. *Application of the field stability evaluation methods to the snow conditions of the eastern Pyrenees*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 386-392.
- Ross, C. and Jamieson, J.B., 2008. *Comparing fracture propagation tests and relating test results to snowpack characteristics*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 376-385.
- Schweizer, J. and Wiesinger, T., 2001. Snow profile interpretation for stability evaluation. *Cold Regions Science and Technology*, 33(2-3): 179-188.
- Schweizer, J., Camponovo, C., Fierz, C. and Föhn, P.M.B., 1995a. Skier triggered slab avalanche release - some practical implications. In: F. Sivardière (Editor), Les apports de la recherche scientifique à la sécurité neige, glace et avalanche. Actes de Colloque, Chamonix, France, 30 mai-3 juin 1995. ANENA, Grenoble, France, pp. 309-315.
- Schweizer, J., Jamieson, J.B. and Schneebeli, M., 2003. Snow avalanche formation. *Rev. Geophys.*, 41(4): 1016, doi:10.1029/2002RG000123.
- Schweizer, J., Schneebeli, M., Fierz, C. and Föhn, P.M.B., 1995b. Snow mechanics and avalanche formation: Field experiments on the dynamic response of the snow cover. *Surveys in Geophysics*, 16(5-6), 621-633.

Continued on page 32 ►

SAN JUAN HISTORY

continued from previous page

A world of uncertainties

Friedman asked what kept my interest in the job. Probably a few things. The winding asphalt ribbon of Highway 550 draws you in with its beauty and diversity of topography, altitudes changes, and the resulting storm conditions. The road climbs out of the sub-alpine zone of Ouray at about 7811' and crests in the alpine on the top of Red at 11,075', then drops into Silverton at 9,305', and climbs again to Molas/Coal Bank at 10,400', dropping quickly into the high desert of Durango. But more importantly, the people I work with are the biggest bonus by far. The real road warriors like Noel Peterson, Ted Vickers, Gary King, Greg Roth, and Dennis McCoy, to name just a few, have made the job very interesting and enjoyable. On each shoot, the protocol has always been safety-first followed with camaraderie and enjoyment. The history and myth of the "old-timers" are a constant source of entertainment and learning. Noel Peterson told me with a smile that the CDOT slogan was always, "Never let the truth get in the way of a good story." Some fine memories and tall tales have come from the CDOT crews.

At times you can feel pretty isolated on the high wire, an elevated and risky autonomy. Pressure builds with a big storm, anticipating the paths that will probably hit the road because you've seen the movie before. It's anybody's guess why forecasters do this job. It could be the smell of powder, throwing 50 pound shots from the helicopter, watching hard slab failure release energy over several alpine basins at once, or maybe just the company you keep.


Whatever the reasons, you get hooked on the excitement and the challenges of the job. It requires a lot of field experience (series of non-fatal errors), collection of empirical evidence, listening to your inner voice (intuition), and distilling all of the variables to reduce uncertainties until you can finally make a decision that you can live with. There are many truths to be learned. It's no big mystery; you pay attention and do your work because you don't want to be a victim of your own bad planning. It helps to be comfortable in the world of uncertainties.

My involvement with the Eddie Imel/Danny Jaramillo tragedy hit me with some harsh realities. A friend of mine calls Red Mountain Pass, "The Living Highway," and I finally realized how accurate his description was. Merely driving the road in winter is a challenge. Add storms, avalanches, rock/ice fall, and night driving to the equation, and it adds up quickly to a difficult and potentially hazardous journey. On a good day there are risks and a bad day can be adventurous.

Jerry Roberts prowls Red Mountain Pass, where he tracks the cast of usual suspects above and below the snow surface. See TAR 27/3 for Part 1 of Jerry Roberts' avalanche history of the San Juan mountains. ❄️



Batman had the bat phone for crime fighting, but Jerry has the duck phone for avalanche fighting.



Low-tech but effective, bore-sighting the 105 howitzer. Photo by Ann Mellick



Two bikers negotiate through Telescope slide debris that ran naturally. Photo by Andy Gleason



A crack following a whumpf on a slope in southwest Montana. Photo by Karl Birkeland

FRACTURE PROPENSITY

continued from pg 29

- Schweizer, J., McCammon, I. and Jamieson, J.B., 2006. *Snow stability evaluation using concepts of fracture mechanics*. In: A. Gleason (Editor), Proceedings of the 2006 International Snow Science Workshop, Telluride, Colorado, pp. 211-218.
- Seligman, G., 1936. Snow structure and ski fields. *International Glaciological Society*, Cambridge, 555 pp.
- Sigrist, C., 2006. *Measurement of fracture mechanical properties of snow and application to dry snow slab avalanche release*. Ph.D. Thesis, Swiss Federal Institute of Technology, Zurich. 139 pp.
- Sigrist, C. and Schweizer, J., 2007. Critical energy release rates of weak snowpack layers determined in field experiments. *Geophysical Research Letters*, 34(3): L03502.
- Simenhois, R. and Birkeland, K.W., 2008a. *The effect of changing slab thickness on fracture propagation*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 755-760.
- Simenhois, R. and Birkeland, K.W., 2008b. *Extended column test results during surface warming: Does surface warming affect fracture propagation propensity?* In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 26-35.
- van Herwijnen, A. and Jamieson, J.B., 2002. *Interpreting fracture character in stability tests*, Proceedings ISSW 2002, International Snow Science Workshop, Penticton BC, Canada, 29 September-4 October 2002, pp. 514-522.
- van Herwijnen, A. and Jamieson, J.B., 2004. *Fracture character in compression tests*. In: K. Elder (Editor), International Snow Science Workshop, Jackson Hole, Wyoming, pp. 182-191.
- van Herwijnen, A. and Jamieson, J.B., 2005. High speed photography of fractures in weak layers. *Cold Regions Science and Technology*, 43(1-2): 71-82.
- van Herwijnen, A., Heierli, J. and Schweizer, J., 2008. *Field study on fracture propagation in weak snowpack layers*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings ISSW 2008, International Snow Science Workshop, Whistler, Canada, 21-27 September 2008, pp. 1-8.
- Winkler, K. and Schweizer, J., 2008. *Comparison of different snow stability tests including the extended column test*. In: C. Campbell, S. Conger and P. Haegeli (Editors), Proceedings of the 2008 International Snow Science Workshop, Whistler, B.C., pp. 393-400.

ACKNOWLEDGMENTS

Many colleagues provided excellent and useful discussions on the topics in this article.

While not a comprehensive list by any means, we would like to thank Dave Gauthier, Joachim Heierli, Alec van Herwijnen, Kalle Kronholm, Ian McCammon, Christian Sigrist, and Ron Simenhois. In addition, Ron Simenhois and Ian McCammon reviewed this paper and provided useful comments.



Karl Birkeland (center) is an avalanche scientist with the USDA Forest Service National Avalanche Center in Bozeman, Montana. Between snowpits he is always hoping to take a few powder turns, and he especially likes chasing his daughters around Bridger Bowl in the winter.

Jürg Schweizer (right) is a senior research scientist and head of the Avalanche Formation group at SLF Davos, Switzerland. He continues to be a great fan of the grandfather of most snowpack tests: the rutschblock. However, he has to admit that the snow saw has become his favorite tool for playing in the snow.

Bruce Jamieson (left) holds the NSERC Research Chair in Snow Avalanche Risk Control at University of Calgary, and supervises the ASARC field program. He is rather fond of digging square holes in the snow and watching how snow fractures. ❄️